

Searching for X-ray Variability in the Glitching Anomalous X-ray Pulsar 1E 1841–045 in Kes 73

Weiwei Zhu¹, Victoria M. Kaspi^{1,2},

ABSTRACT

Anomalous X-ray Pulsars (AXPs) are now established to exhibit significant X-ray variability and be prolific glitchers, with some glitches being accompanied by large radiative changes. An open issue is whether AXP glitches are generically accompanied by radiative changes, relevant for understanding magnetar physical properties. Here we report on an analysis of archival X-ray data from the AXP 1E 1841–045, obtained between 1993 and 2007. This AXP, located in the center of SNR Kes 73, has exhibited three glitches between 2002 and 2007, as determined by *RXTE* monitoring since 1999. We have searched for evidence of phase-averaged flux variability that could be present if glitches in AXPs are usually accompanied by radiative changes. We find no evidence for glitch-correlated flux changes from this source, arguing that such behavior is not generic to AXPs.

Subject headings: pulsars: individual (1E 1841–045) — X-rays: stars — stars: neutron

1. Introduction

X-ray variability in AXPs is now established as a characteristic property of this source class. Their fluxes and spectra vary on a variety of different time scales, from a few milliseconds (e.g. Gavril et al. 2002), to several years (Dib et al. 2007). See Kaspi (2007); Mereghetti (2008) for recent reviews of AXP variability.

AXPs are also now known to be prolific glitchers, and include some of the most active glitchers known in the neutron-star population (Dib et al. 2008). By ‘glitch’ we mean a sudden rotational spin-up, or similar timing anomaly. One of the largest glitches seen so far, from AXP 1E 2259+586, was accompanied by major radiative changes, including bursts and

¹Department of Physics, McGill University, Montreal, QC, H3A 2T8, Canada; zhuww@physics.mcgill.ca, vkaspi@physics.mcgill.ca

²Canada Research Chair; Lorne Trottier Chair; R. Howard Webster Fellow of CIFAR

a factor of ~ 20 pulsed and persistent flux increase (Kaspi et al. 2003; Woods et al. 2004). Dib et al. (2009) showed that glitches in AXP 1E 1048.1–5937 were also accompanied with radiative enhancements. Such events are thought to be the result of sudden yielding of the neutron-star crust due to internal stresses caused by the decay of the magnetar-strength field. The restructuring results in changes to the stellar interior – as evidenced by the glitch – and to the stellar exterior – as evidenced by the dramatic radiative changes. Observations of such AXP outbursts are a potentially powerful probe of the physics of magnetars (e.g. Eichler & Shaisultanov 2010).

Links between the X-ray variability and glitches of AXP RXS J170849.0–400910 have also been reported (Rea et al. 2005; Campana et al. 2007; Israel et al. 2007; Götz et al. 2007). These authors suggest the existence of a general correlation between magnetars’ flux and glitch epochs. Specifically, Götz et al. (2007) reported $\sim 40\%$ flux changes in RXS J170849.0–400910 based on eight observations made over the course of ~ 9 yr, during which they report four glitches. If correct, this suggests that glitches are usually, and possibly always, accompanied with radiative changes. However, the sparsity of observing epochs compared to the number of glitches for this AXP thus far is problematic in proving the variability is glitch-correlated. Additionally, no comparable pulsed flux changes were observed in the same time span, during which such measurements were available regularly on a monthly basis (Dib et al. 2008). This apparent conflict could, however, be explained if the pulsed fraction were precisely anti-correlated with flux. An anti-correlation between pulsed fraction and flux has been seen in AXP 1E 1048.1–5937 (Tiengo et al. 2005; Tam et al. 2008), although not to a degree that render pulsed flux variations absent.

Here we investigate the hypothesis that AXP glitches are generically accompanied by radiative changes by considering AXP 1E 1841–045 in supernova remnant (SNR) Kes 73, one of the most frequent glitchers among AXPs. From *RXTE* monitoring, we know that 1E 1841–045 has had three glitches between 1999 and 2008 (Dib et al. 2008). The glitches occurred on 2002 July 9, 2003 December 24, and 2006 March 29, and had $\Delta\nu/\nu$ of 5.63×10^{-6} , 2.45×10^{-6} and 1.39×10^{-7} , respectively, and were not accompanied by any X-ray pulsed flux changes. If, as seen in 1E 2259+586 and 1E 1048.1–5937 and reported for RXS J170849.0–400910, glitches are generically accompanied by radiative changes, and, if as for RXS J170849.0–400910, such radiative changes are not necessarily apparent in the pulsed flux data, it is possible that the phase-averaged flux (unavailable from *RXTE* monitoring) varies in concert with glitches in 1E 1841–045. We investigate this possibility here. Also worth noting is that Gotthelf et al. (1999) studied the timing of AXP 1E 1841–045 using archival *GINGA*, *ASCA*, *ROSAT* and *RXTE* data taken between 1993 and 1999 and found no glitches with $\Delta\nu/\nu > 5 \times 10^{-6}$.

In this paper, we report on our analysis of archival X-ray data for 1E 1841–045 collected by *ASCA*, *Chandra*, *XMM* and *Suzaku* during the past 17 years, including two observations that were made fortuitously very closely following glitches. We have looked for correlations between the AXP’s flux variability and glitch epochs. In section 2 we describe the observations and data reduction process. In section 3 we describe our spectral analysis and AXP flux extraction method. Our results and conclusions are discussed in section 4.

2. Observations

For this study, we searched online X-ray archives for all existing observations of the Kes 73 field. We found a total of eleven observations from four different focussing X-ray observatories (listed in Table 1). We do not include in our analysis the many *RXTE* observations, as due to its non-focussing nature, these provide only pulsed flux measurements, and are already published (Dib et al. 2008). Next we report on our analysis of the eleven focussing-telescope observations.

2.1. *ASCA* Observations

Seven *ASCA* (Tanaka et al. 1994) observations of the AXP 1E 1841–045 and the SNR Kes 73 were taken between 1993 and 1999 (see Table 1). Our analysis began with the screened data from the two Gas Imaging Spectrometers (GISs, Burke et al. 1994), which were filtered with the standard revision 2 screening criteria. Given the angular resolution of the GISs, the SNR ($\sim 2'$) was unresolvable in the images, therefore the spectra we extracted from the GISs contain photons from both the AXP and the SNR. Using the `ftool xselect`¹, we extracted spectra from source regions of $9'.8$ radius (a region large enough to encircle the extended emission from AXP 1E 1841–045 and Kes 73) and background spectra from regions of radius $\sim 5'$, away from the source region for all the *ASCA* GIS observations. The GIS spectra were then combined with Redistribution Matrices File (RMF) of the GISs and Auxiliary Response File (ARF) files generated using the `ftool ascaarf`, and grouped with a minimum of 25 counts per bin. Finally the exposure of the grouped spectra were corrected for the deadtime effect using the `ftool deadtime`. In this study, we did not include the spectra extracted from the two Solid-state Imaging Spectrometers (SISs, Ohashi 1996), primarily because of the significantly fewer counts collected by these instruments.

¹<http://heasarc.gsfc.nasa.gov/ftools/xselect/>

2.2. *Chandra* Observation

The AXP and SNR were observed by the *Chandra X-ray Observatory* with the Advanced CCD Imaging Spectrometer (ACIS, Garmire et al. 2003) in timed exposure (TE) mode on 2000 July 23 and in continuous clocking (CC) mode on 2000 July 29 (Table 1). The data were analyzed and reported by Morii et al. (2003). Here we used *Ciao* version 3.4². Because the spectrum of the pulsar in TE mode was heavily affected by pile-up, we did not use these data; instead we extracted the spectrum of AXP 1E 1841–045 from the CC mode data. In these data, the image of the pulsar and SNR were collapsed into one dimension. From the level 2 event list provided by the *Chandra* X-ray Center, we extracted the spectrum of the pulsar using a box-shaped region capturing a 2".5 long segment along the one-dimensional image and centered on the pulsar. The background spectrum was extracted from two 7".5 long segments adjacent to the source region. The resulting source and background spectra were then combined with RMF and ARF files generated using the `psextract` command and grouped with a minimum of 25 counts per bin.

2.3. *XMM* Observations

The AXP was observed by the *Newton X-ray Multi-Mirror Mission (XMM)* observatory (Jansen et al. 2001) on 2002 October 5 and 2002 October 7 (Table 1) with the European Photon Imaging Camera (EPIC) pn (Strüder et al. 2001) camera operating in large window mode and the EPIC MOS cameras (Turner et al. 2001) in full window mode. For our analysis, we used the *XMM* Science Analysis System (**SAS**) version 8.0.0³ and calibrations (updated 2008 Oct 3). Given the pn and MOS cameras' angular resolution, 1E 1841–045 can be resolved from Kes 73. For the two *XMM* observations, we used only the data from the EPIC pn camera to take advantage of its larger photon collecting area and to avoid cross-calibration issues between the pn camera and the mos cameras. We extracted the pulsar's spectrum from a circular region of radius 32".5 (a radius large enough to capture more than 90% of the photon events from the point source) centered on the pulsar. Background spectra were extracted from an annular region of radius between 35" and 115" centered on the pulsar. The pulsar spectrum was grouped with a minimum of 25 counts per bin and then combined with the background spectrum and RMF and ARF files generated by the **SAS** software.

²<http://131.142.185.90/ciao3.4/index.html>

³See <http://xmm.esac.esa.int/sas/8.0.0/>

2.4. *Suzaku* Observations

AXP 1E 1841–045 and Kes 73 were also observed by the *Suzaku* observatory (Mitsuda 2007) on 2006 April 19 (Table 1). The data analysis was reported by Morii et al. (2008). Onboard *Suzaku*, there are two X-ray detectors: the X-ray Imaging Spectrometer (XIS, Koyama 2007, consisting of four CCDs sensitive in soft-X-ray band) and the Hard X-ray Detector (HXD, Takahashi 2007, sensitive to 10–600 keV X-rays). Here we present a spectral analysis of the XIS data only. Given the angular resolution of the XIS, the SNR was unresolvable in the XIS image. Therefore the spectra we extracted from the XIS detectors contain photons from both the AXP and the SNR. We used cleaned events screened by the standard pipeline processing version 2.0.6.13⁴. The source spectra were extracted from a circular region of 260'' radius. Background spectra were extracted from an annulus region of radius between 260'' and 520''. The extracted spectra were grouped with a minimum of 25 counts per bin, and then combined with the RMF and ARF files generated using the `ftools` `xisrmfgen` and `xissimarfgen`.

3. Spectroscopy

Among the eleven observations we used, the AXP can be resolved from the SNR Kes 73 only in the *Chandra* and *XMM* observations. For these, it is possible to extract either the neutron star’s spectrum or the combined spectrum of the neutron star and the SNR. By contrast, only combined spectra can be extracted from the *ASCA* and *Suzaku* observations.

The spectra of AXPs are often parametrized by a blackbody plus a power-law (although this is known to be an approximation to a likely Comptonized blackbody spectrum – see Thompson et al. 2002 and, for example, Rea et al. 2008). The spectra of SNRs are often fit with models like, for example, VSEDOV (a plane-parallel shock radiation model with separate ion and electron temperatures), VNEI (a non-equilibrium ionization collisional plasma model), or VPSHOCK (a plane-parallel shocked plasma model). See Borkowski et al. 2001 for a review of these models. In this paper, we modeled the neutron star radiation with a blackbody plus power-law (BB+POW), and the SNR radiation with a VSEDOV model, using `xspec` 12.5.0⁵. The focus of our investigation is on the AXP’s X-ray flux; in modelling the SNR we sought only a suitable parameterization to allow us to subtract off its flux reliably. As we show below, the VSEDOV model is adequate for these purposes.

⁴http://heasarc.nasa.gov/docs/suzaku/processing/criteria_xis.html

⁵<http://heasarc.nasa.gov/docs/xanadu/xspec/>

We modeled the interstellar absorption by multiplying a WABS (a photo-electric absorption model, in which the interstellar absorption is characterized by a single parameter N_H , the neutral hydrogen column density along the line of sight) model to both the BB+POW and VSEDOV models.

Figure 1 shows the combined spectra from *XMM* and the components of the best-fit model. The AXP power-law component clearly dominates the spectra above ~ 4 keV. Therefore, we chose to study the AXP flux in the 4–10 keV band only, in order to minimize SNR contamination. Nevertheless, we still attempted to remove the remaining small contribution of SNR in this band for the fluxes measured from *ASCA* and *Suzaku* observations, so that we could compare them with the neutron-star-only fluxes measured with *Chandra* and *XMM*.

It is reasonable to assume that both the interstellar absorption and SNR radiation do not change over a time scale of about a decade. Consequently, in our attempt to remove the SNR flux, we used the same N_H and VSEDOV parameters for all the spectra of the different observations, and allowed only the BB+POW model to vary from observation to observation.

When fitting a WABS(BB+POW+VSEDOV) model to those spectra containing emission from both the AXP and SNR, it is challenging to constrain the normalization parameters of both the BB and VSEDOV models simultaneously, because these two models dominate the same energy band and their parameters are highly covariant. Fortunately, the neutron star can be spatially resolved out in the *XMM* observations, so we can use them to determine the relative strength of the two spectral components. We therefore fit a WABS(BB+POW+VSEDOV) model to the combined (AXP+SNR) *XMM* spectrum. To ensure that we had the correct BB+POW model for the neutron star, we simultaneously fit the spectrum extracted from only the neutron star, requiring common neutron-star spectral parameters. Hence we could determine the parameters and normalization of the VSEDOV model for the SNR.

To further improve the spectral model, next we included the *ASCA* and *Suzaku* spectra and the *Chandra* CC-model neutron-star spectrum and performed a large joint fit, which required multiple iterations in order to converge. The result was a value for N_H and for the VSEDOV model parameters that fit all the SNR spectra reasonably well, albeit not perfectly (reduced $\chi^2 = 1.36$ for 3801 degrees of freedom for the joint fit). The best-fit N_H and VSEDOV parameters can be found in Table 2.

Note that for all the spectral fitting described above, we used the 0.8–10 keV band. However for *Suzaku*, we found that there were always significant residuals in the range 1.7–3.5 keV and above 9 keV, and in general, these residuals differed significantly among the

four XIS instruments. Therefore, we ignored the 1.7–3.5 keV band and above 9 keV for the *Suzaku* spectra. Furthermore, the SNR Kes 73 is larger than the field-of-view of the *Suzaku* XISs and was not entirely captured. Therefore, when fitting the *Suzaku* spectrum, we allowed the VSED OV normalization parameter to vary.

Finally, by using the best-fit N_H and VSED OV model, we could remove the SNR flux contribution from the *ASCA* and *Suzaku* observations in separate fits to their spectra, hence measuring the AXP fluxes. For the *Chandra* and *XMM* observations, we simply fitted the resolved AXP spectra with an absorbed BB+POW model. The best-fit BB+POW parameters and the measured 4–10 keV unabsorbed fluxes of the AXP are presented in Table 3. The fluxes are also plotted in Figure 2. The uncertainties we report on the 4–10 keV unabsorbed AXP fluxes were estimated by the measured fractional uncertainties on the 4–10 keV absorbed total fluxes. For those observations in which the best-fit reduced χ^2 was larger than unity, we multiplied the reported flux uncertainties by the square root of the reduced χ^2 . This is to account for systematic errors in our imperfect modelling of the SNR spectrum.

To confirm the robustness of the measured AXP flux values, we repeated the entire above analysis by modelling the SNR spectra in different ways. We found that both the VNEI and VPSHOCK models can fit the SNR spectra as well as the VSED OV model, and that the 4–10 keV unabsorbed neutron star fluxes we measured using these models are consistent within reported uncertainties with the values we found using the VSED OV model. Therefore, we feel confident that the phase-averaged AXP flux values we report are well constrained and robust. However, the BB+POW spectral parameters we measured when fitting the SNR with different models were not as robust as the 4–10 keV fluxes, changing significantly with SNR model. As we do not believe them to be reliably determined, we do not quote their uncertainties in Table 3, and do not consider them further.

Also in Figure 2, we show the AXP’s 2–10 keV pulsed flux as measured in monitoring observations with *RXTE* since early 1999. Details about how these pulsed fluxes were determined are provided in Dib et al. (2008). The *RXTE* pulsed fluxes show no significant variations. The vertical lines in Figure 2 indicate the epochs of the three observed glitches (Dib et al. 2008).

4. Discussion and Summary

The goal of this study was to see whether the prolific glitching AXP 1E 1841–045 shows phase-averaged flux variability, in spite of showing no evidence for pulsed flux variability. Also, we wished to determine whether any variability is correlated with its glitches as has

been seen in AXP 1E 2259+586 in its 2002 major outburst, in 1E 1048.1–5937, and also reported for RXS J170849.0–400910.

As is clear from Panel a of Figure 2, in the 4–10 keV band, the neutron star’s flux did not vary by more than $\sim 30\%$ in 13 years. Interestingly the largest variations we find are in the multiple pre-1999 *ASCA* observations: in those seven observations, a fit to a constant flux results in a reduced χ^2 of 3.7 for 6 degrees of freedom, which has a probability of occurring by chance of ~ 0.001 . However, during this time, there were certainly no large glitches ($\Delta\nu/\nu < 5 \times 10^{-6}$) (Gotthelf et al. 1999).

The *ASCA* fluxes all appear to be higher than the fluxes measured from other observations. If this difference is accurate, then the AXP’s flux must have dropped around 2000, and we might in principle expect some change in the pulsed flux as monitored by *RXTE* at that epoch. However, the pulsed flux of the pulsar was constant around that time as seen in Panel b of Figure 2. Therefore, we suspect that the relative increase in the flux in the *ASCA* observations compared with those of the other observatories may be due to instrumental calibration issues.

The fluxes measured from the last four observations taken by *Chandra*, *XMM* and *Suzaku* can be fitted with a constant flux model (reduced $\chi^2 = 1.9$ for 3 degrees of freedom, corresponding to a probability of having occurred by chance of 0.125). Thus we conclude that the phase-averaged 4–10 keV fluxes of the last four observations were consistent with being constant, and we put an upper limit of 11% on long-term variability in this energy band. Importantly, the two *XMM* observations were taken only 88 and 90 days after the first glitch, and the *Suzaku* observation was taken only 27 days after the third glitch. By contrast, the 4–10 keV flux of 1E 2259+586 was 50% higher than in quiescence 21 days after its 2002 glitch (Zhu et al. 2008), that of 1E 1048.1–5937 was a factor of 6 higher ≥ 38 days after its 2007 glitches (Tam et al. 2008), and was 50–70% higher for RXS J170849.0–400910 ~ 53 days after its first 2005 glitch as inferred from Götz et al. (2007)⁶. Therefore, we conclude that unlike in 1E 2259+586, 1E 1048.1–5937 and possibly RXS J170849.0–400910, there is no evidence for glitch-correlated flux changes in AXP 1E 1841–045.

One caveat of our study is that we were limited to the harder part of the neutron star’s emission spectrum. The flux from the blackbody component was not well constrained. Therefore, we cannot rule out changes in the neutron star’s thermal radiation, only changes

⁶This is calculated based on the reported 1–10 keV fluxes of RXS J170849.0–400910 from its 2003 *XMM* observation and 2005 *Swift* observation. We assumed a power-law to blackbody flux ratio of 3 for the 1–10 keV fluxes, and then calculated the 4–10 keV fluxes using webPIMMS (<http://heasarc.gsfc.nasa.gov/Tools/w3pimms.html>)

in the power-law component in the 4–10 keV band, which we note constitutes ~ 0.25 of the stellar flux (BB+POW) in the 1–10 keV band.

Thus our results argue against the hypothesis that there exists a generic correlation between the X-ray flux variability and glitch epochs in AXPs, further supporting the argument that glitches in AXPs can be either radiatively loud or radiatively silent (Dib et al. 2008). There is of course precedent for radiatively silent glitches in neutron stars, in that no rotation-powered pulsar glitch has ever been reported to be accompanied with any radiative change, although rapid X-ray follow-up has only been accomplished in one case (Helfand et al. 2001). Any physical model of magnetar glitches will have to explain the simultaneous existence of both types. This is true of even a single source, as there is evidence that AXP 1E 2259+586 has both, given that its most recent glitch showed no pulsed flux change (Dib et al. 2008b). Recently Eichler & Shaisultanov (2010) have argued that AXP glitches are triggered by energy releases at depths below ~ 100 m in the crust, with angular momentum vortex unpinning being due to global mechanical motion triggered by the energy release, not by heat as has been proposed in the context of rotation-powered pulsars glitches (Link & Epstein 1996; Link & Cutler 2002). If mechanical triggering occurs, then radiatively silent glitches of the same amplitude as radiatively loud glitches are possible since less energy is required to trigger a glitch than to cause a substantial X-ray brightening. If so, then Eichler & Shaisultanov (2010) predict that all AXP glitches should occur simultaneously with or before the observed X-ray brightening; this can be tested with continuous (daily or better) X-ray monitoring observations. Moreover, although mechanical unpinning of vortices by activity in the lower crust does result in heat release, the latter could take as much as several years to reach the surface. This could help explain the long-term X-ray variability trends that have been reported in some AXPs (e.g. Dib et al. 2007).

We thank Koji Mukai of the *ASCA* GOF for assistance with the *ASCA* data and David Eichler for useful conversations. We also thank Rim Dib for providing the *RXTE* pulsed fluxes. VMK acknowledges support from NSERC via a Discovery Grant, FQRNT via the Centre de Recherche en Astrophysique du Québec, CIFAR and holds a Canada Research Chair and Lorne Trottier Chair in Astrophysics and Cosmology.

REFERENCES

- Borkowski, K. J., Lyerly, W. J., & Reynolds, S. P. 2001, *ApJ*, 548, 820
- Burke, B. E., Mountain, R. W., Daniels, P. J., Cooper, M. J., & Dolat, V. S. 1994, *IEEE Transactions on Nuclear Science*, 41, 375

- Campana, S., Rea, N., Israel, G. L., Turolla, R., & Zane, S. 2007, *A&A*, 463, 1047
- Dib, R., Kaspi, V. M., & Gavriil, F. P. 2007, *ApJ*, 666, 1152
- . 2008, *ApJ*, 673, 1044
- . 2008b, in *American Institute of Physics Conference Series*, Vol. 983, 40 Years of Pulsars: Millisecond Pulsars, Magnetars and More, ed. C. Bassa, Z. Wang, A. Cumming, & V. M. Kaspi, 262
- . 2009, *ApJ*, 702, 614
- Garmire, G. P., Bautz, M. W., Ford, P. G., Nousek, J. A., & Ricker, G. R. 2003, *Proc. SPIE*, 4851, 28
- Gavriil, F. P., Kaspi, V. M., & Woods, P. M. 2002, *Nature*, 419, 142
- Gotthelf, E. V., Vasisht, G., & Dotani, T. 1999, *ApJ*, 522, L49
- Götz, D., Rea, N., Israel, G. L., Zane, S., Esposito, P., Gotthelf, E. V., Mereghetti, S., Tiengo, A., & Turolla, R. 2007, *A&A*, 475, 317
- Eichler, D., Shaisultanov, R. 2010, *ApJ*, submitted
- Helfand, D. J., Gotthelf, E. V., & Halpern, J. P. 2001, *ApJ*, 556, 380
- Israel, G. L., Götz, D., Zane, S., Dall’Osso, S., Rea, N., & Stella, L. 2007, *A&A*, 476, L9
- Jahoda, K., Swank, J. H., Giles, A. B., Stark, M. J., Strohmayer, T., Zhang, W., & Morgan, E. H. 1996, *Proc. SPIE*, 2808, 59
- Jansen, F. et al. 2001, *A&A*, 365, L1
- Link, B., Epstein, R. I. 1996, *ApJ*, 457, 844
- Link, B., Cutler, C. 2002, *MNRAS*, 336, 211
- Kaspi, V. M. 2007, *Ap&SS*, 308, 1
- Kaspi, V. M., Gavriil, F. P., Woods, P. M., Jensen, J. B., Roberts, M. S. E., & Chakrabarty, D. 2003, *ApJ*, 588, L93
- Koyama, K. e. a. 2007, *PASJ*, 59, 23
- Link, B. & Cutler, C. 2002, *MNRAS*, 336, 211

- Mereghetti, S. 2008, *A&A Rev.*, 15, 225
- Mitsuda, K. e. a. 2007, *PASJ*, 59, 1
- Morii, M. et al. 2008, in *American Institute of Physics Conference Series*, Vol. 983, 40
Years of Pulsars: Millisecond Pulsars, Magnetars and More, ed. C. Bassa, Z. Wang,
A. Cumming, & V. M. Kaspi, 268–270
- Morii, M., Sato, R., Kataoka, J., & Kawai, N. 2003, *PASJ*, 55, L45
- Ohashi, T. e. a. 1996, *PASJ*, 48, 157
- Rea, N., Oosterbroek, T., Zane, S., Turolla, R., Méndez, M., Israel, G. L., Stella, L., &
Haberl, F. 2005, *MNRAS*, 361, 710
- Rea, N., Zane, S., Turolla, R., Lyutikov, M., Götz, D. 2008, *ApJ*, 686, 1245
- Strüder, L. et al. 2001, *A&A*, 365, L18
- Takahashi, T. e. a. 2007, *PASJ*, 59, 35
- Tam, C. R., Gavriil, F. P., Dib, R., Kaspi, V. M., Woods, P. M., & Bassa, C. 2008, *ApJ*,
677, 503
- Tanaka, Y., Inoue, H., & Holt, S. S. 1994, *PASJ*, 46, L37
- Thompson, C., Lyutikov, M., & Kulkarni, S. R. 2002, *ApJ*, 574, 332
- Tiego, A. and Mereghetti, S. and Turolla, R. and Zane, S. and Rea, N. and Stella, L. and
Israel, G. L. 2005, *A&A*, 437, 997
- Turner, M. J. L. et al. 2001, *A&A*, 365, L27
- Woods, P. M., Kaspi, V. M., Thompson, C., Gavriil, F. P., Marshall, H. L., Chakrabarty,
D., Flanagan, K., Heyl, J., & Hernquist, L. 2004, *ApJ*, 605, 378
- Zhu, W., Kaspi, V. M., Dib, R., Woods, P. M., Gavriil, F. P., & Archibald, A. M. 2008,
ApJ, 686, 520

Table 1. X-ray observations of AXP 1E 1841–045 used in this study.

Date	Observatory	t^a (ks)	Offset ^b (')
1993 Oct 11	<i>ASCA</i>	40	6.8
1997 Apr 21	<i>ASCA</i>	9	7.3
1998 Mar 27	<i>ASCA</i>	39	6.5
1999 Mar 22	<i>ASCA</i>	20	5.3
1999 Mar 29	<i>ASCA</i>	20	5.2
1999 Apr 06	<i>ASCA</i>	19	5.2
1999 Apr 13	<i>ASCA</i>	21	5.2
2000 Jul 29	<i>Chandra</i>	10	0.097
2002 Oct 05	<i>XMM</i>	3.8	1.152
2002 Oct 07	<i>XMM</i>	4.4	1.144
2006 Apr 19	<i>Suzaku</i>	98	3.9

^aThe effective exposure time of the instrument used for the spectral analysis in this paper.

^bThe pointing offsets of the observations relative to the position of the AXP.

Table 2. WABS and VSED OV parameter values

Parameter	Value
N_H	$2.77 \times 10^{22} \text{cm}^{-2}$
kT_a	0.56 keV
kT_b	0.56 keV
Mg	0.63 ^a
Si	1.10
S	1.92
Ca	0.32
Fe	0.57
Ni	2.51
τ	$1.50 \times 10^{11} \text{ s}$
Redshift	0.00
Norm ^b	0.33
$H...Ar^c$	1.0

^aThe element abundances quoted here are the relative abundances based on the Solar mixture abundances.

^bNormalization parameter for the VSED OV model, $\frac{10^{-14}}{4\pi[D_A(1+z)]^2 \int n_e n_H dV}$ where D_A is the angular diameter distance to the source (cm), and n_e, n_H (cm^{-3}) are the electron and hydrogen densities respectively.

^cElements $H...Ar$: $H He C N O Ne$ and Ar were fixed to the solar abundance because the SNR spectra are not sensitive to them.

Table 3. Measured AXP model parameters and fluxes.

Date	Observatory	MJD	kT^a (keV)	Γ^a	F_{NS}^b ($10^{-11}\text{erg cm}^{-2}\text{s}^{-1}$)
1993 Oct 12	<i>ASCA</i>	49272.1	0.20	2.72	1.16(4)
1997 Apr 21	<i>ASCA</i>	50559.6	0.19	3.00	1.2(1)
1998 Mar 27	<i>ASCA</i>	50899.95	0.20	2.70	1.36(5)
1999 Mar 22	<i>ASCA</i>	51259.7	0.20	2.89	1.29(4)
1999 Mar 30	<i>ASCA</i>	51267.0	0.19	2.80	1.39(5)
1999 Apr 06	<i>ASCA</i>	51274.6	0.21	2.74	1.39(5)
1999 Apr 14	<i>ASCA</i>	51282.3	0.21	2.68	1.36(4)
2000 Jul 29	<i>Chandra</i>	51754.32	0.40	2.10	0.975(3)
2002 Oct 05	<i>XMM</i>	52552.16	0.22	2.59	1.05(4)
2002 Oct 07	<i>XMM</i>	52554.16	0.21	2.47	1.08(3)
2006 Apr 19	<i>Suzaku</i>	53844.8	0.38	1.94	1.03(2)

^aThe reported best-fit kT and Γ parameters of the BB+POW model vary depending on the assumed SNR model; we do not report the fit uncertainties of these parameters as they do not reflect the true uncertainties. The values, determined while assuming the VSED OV model for SNR, are provided for reference only.

^bPhase-averaged unabsorbed neutron-star flux in the 4–10 keV band. The number in parenthesis represents the 1σ uncertainties in the last digit. See text for details.




Fig. 1.— The unfolded *XMM* pn spectra and components of the best-fit BB+POW+VSED OV model.




Fig. 2.— (a) 4–10 keV unabsorbed phase-averaged fluxes from AXP 1E 1841–045, as determined by our analysis. (b) 2–10 keV pulsed fluxes from the pulsar as measured by *RXTE*, in units of counts per second per Proportional Counter Unit (PCU) (see Dib et al. 2008, for details).